

First records of *Anthidium florentinum* (Hymenoptera, Megachilidae) in the Canary Islands: insights on citizen science and exotic bee species

Diego Patiño-Sauma¹, Carlos Ruiz¹, John S. Ascher²

1 Department of Animal Biology, Edaphology and Geology, University of La Laguna, 38206 La Laguna, Tenerife, Canary Islands, Spain **2** Department of Biological Sciences, National University of Singapore, 16 Science Drive 4, Singapore 117558, Singapore

Corresponding author: Diego Patiño-Sauma (diegopsauma@gmail.com)

Academic editor: Christopher K. Starr | Received 27 September 2024 | Accepted 16 December 2024 | Published 5 February 2025

<https://zoobank.org/2BFAD9B2-8763-4300-B375-9E9EDE170084>

Citation: Patiño-Sauma D, Ruiz C, Ascher JS (2025) First records of *Anthidium florentinum* (Hymenoptera, Megachilidae) in the Canary Islands: insights on citizen science and exotic bee species. Journal of Hymenoptera Research 98: 85–93. <https://doi.org/10.3897/jhr.98.137948>

Abstract

In recent years, the development of biodiversity inventorying platforms has contributed to the detection of exotic species, bringing awareness to the general public about their impact on native habitats. This is important in the Canary Islands and other oceanic archipelagos, where invasive species pose a great risk to the conservation of ecosystems and biodiversity monitoring strategies may not be completely adequate in their detection. Here we report for the first time the occurrence of the Florentine wool carder bee, *Anthidium florentinum*, in the Canary Islands and address its current distribution in the islands of Tenerife and Gran Canaria with the help of citizen science. Other bee species recently reported as new to the archipelago through the use of iNaturalist are summarized. The possible effects of *A. florentinum* on native bee communities in the archipelago, along with the role of cities in hosting biological invasions are also discussed.

Keywords

Alien species, early detections, oceanic islands, social media, wood-nesting bees

Introduction

Invasive bees are responsible for several negative impacts such as competition for floral and nesting resources, disruption of pollination networks, and introduction of pathogens (Russo 2016). In oceanic islands, native pollinators in most cases have evolved in isolation from the mainland, which makes them especially sensitive to the advent of alien species, owing to a lack of advanced competitive strategies against continental newcomers (Sugiura 2016).

Wood-nesting bee species (particularly many megachilid species from the genera *Anthidium* Fabricius, 1804 and *Megachile* Latreille, 1802) are especially prone to accidental transport by trade, which makes them the most common introduced and invasive species (Russo 2016). In the Canary Islands, several wood-nesting species have been introduced in the last decades, such as the megachilid bees *Anthidium manicatum* (Linnaeus, 1758) (Lieftinck, 1958) and *Megachile otomita* Cresson, 1878 (Strudwick and Jacobi 2018), the carpenter bees *Xylocopa violacea* (Linnaeus, 1758) (Ortiz-Sánchez et al. 2016) and *X. pubescens* Spinola, 1838 (Ruiz et al. 2020), and various *Hylaeus* Fabricius, 1793 species (Lugo et al. 2022).

One wood-nesting species that has also been accidentally introduced is the Florentine wool carder bee *Anthidium florentinum* (Fabricius, 1775). This species has a natural distribution extending from South and Central Europe (Amiet et al. 2004) to China (Niu et al. 2020), and also includes several mediterranean islands such as Malta (Balzan et al. 2017), the Balearic Islands (Baldock et al. 2020), and Cyprus (Varnava et al. 2020). In recent decades, this species has expanded northward beyond its historic range limits (Fig. 1) to reach Switzerland (Kouakou et al. 2008), Slovenia (Gogala 2012), Austria (Zettel et al. 2016), Germany (Schwenninger 2008) and Russia (Antropov et al. 2017) including the Moscow area and a disjunct population as far north as St. Petersburg (reflectitur_photons Obs. <https://www.inaturalist.org/observations/14364056>). *A. florentinum* has also been reported from Algeria (Aguib et al. 2010) and very recently in India (Sardar et al. 2022). Additionally, this species has been recently introduced in North America, as it was observed in eastern Canada from the Montreal area (Normandin et al. 2017) and in the Eastern USA very locally in far Northern New York State just south of the Quebec border (joshualincoln Obs. <https://www.inaturalist.org/observations/93793337>).

A. florentinum is a polylectic and univoltine species with a flying period that starts early in June and ends in late August, overlapping for both males and females (Fortunato et al. 2013). Males carry out a mating system called resource defense polygyny in which they patrol and defend flower patches of different plant families potentially attractive to female foragers, particularly Fabaceae and Lamiaceae (Romero et al. 2021). They show a very aggressive behavior, attacking other conspecific males, and also different species of pollinators, in order to keep intruders away from the area they are securing, to ensure reproductive success (Wirtz et al. 1992).

Anthidium florentinum can be easily mistaken for *A. manicatum*, a common and widespread species that has similar overall appearance, size and colouration, but with




Figure 1. Range of distribution of *Anthidium florentinum* based on georeferenced records (dots) obtained from GBIF 2024 (<https://www.gbif.org/>). Countries within the natural range of distribution of *A. florentinum* are marked with yellow dots. Countries near its native range where this species has been detected in the last decades are marked with blue dots. Regions where *A. florentinum* has been recently introduced are marked with red dots **A** Eurasia and North Africa **B** North America.

subtle differences. For instance, males of both species can be differentiated by looking at the fourth and fifth abdominal tergites, where *A. florentinum* shows a lateral tooth at both sides of these segments, whereas *A. manicatum* only has this trait at tergite 5. Also, *A. manicatum* usually displays a black stain in the upper part of the clypeus in both sexes, whereas *A. florentinum* doesn't. Differences in coloration patterns are useful provided that considerable geographic and individual variation in both species is accounted for.

In this issue we report the occurrence of *A. florentinum* in the islands of Tenerife and Gran Canaria, and discuss the current distribution of this new bee species in the archipelago, as well as the potential threat that this exotic species may represent for native communities of pollinators in the Canary Islands. Additionally, we comment on the role of citizen science in monitoring exotic species, and note its contribution to new discoveries and to a deeper understanding on the spatial distribution of species.

Methods

In May of 2022 two specimens in an urban park of Santa Cruz de Tenerife city were misidentified as *A. manicatum* on iNaturalist (iNaturalist 2024) as it was the only known *Anthidium* in the island. The observation was later identified as *A. florentinum* by one of us (JSA) via iNaturalist later in the year. An active search with nets was performed in different areas of the enclosure where flowering plants were abundant. In addition, we searched for observations submitted by users in social media platforms like Facebook, X, Instagram and iNaturalist (<https://www.inaturalist.org>) to have a broader view of the distribution of *A. florentinum* in the Canary Islands. For the same purpose, the bee collection of the Zoology Department of the University of La Laguna was examined.

Results

Several specimens of *A. florentinum* were either collected by net or uploaded as observations in iNaturalist by users in different localities on the islands of Tenerife and Gran Canaria.

Collected specimens

Spain. Canary Islands, **Tenerife**, Santa Cruz de Tenerife. 1 ♂; Parque García-Sanabria; 28.4715, -16.2543; 12 Jun. 2022; D. Patiño leg. • 3 ♂♂, 3 ♀♀; Parque García-Sanabria; 28.4715, -16.2543; 12 Aug. 2022; D. Patiño leg.

Records obtained from citizen science

Spain. Canary Islands, **Tenerife**. 1 ♂, 1 ♀; Santa Cruz de Tenerife; 28.4729, -16.2543; 30 May 2022; D. Patiño, pers. obs. (<https://www.inaturalist.org/observations/119402491>) • 1 ♀; Arafo; 28.3428, -16.4353; 29 Jun. 2022; Gustavo Peña, pers. obs. (<https://www.inaturalist.org/observations/125579345>) • 1 ♀; Arafo; 28.3428, -16.4353; 10 Aug. 2022; Gustavo Peña, pers. obs. (<https://www.inaturalist.org/observations/130412080>) • 1 ♀; San Cristóbal de La Laguna; 28.4930, -16.3162; 13 May 2023; carmendgq, pers. obs. (<https://www.inaturalist.org/observations/161777634>) • 1 ♂; Santa Cruz de Tenerife; 28.4521, -16.2564; 27 May 2023; fer_guerra, pers. obs. (<https://www.inaturalist.org/observations/164070022>) • 1 ♂; Santa Cruz de Tenerife; 28.4518, -16.2564; 27 May 2023; fer_guerra, pers. obs. (<https://www.inaturalist.org/observations/164070949>) • 1 ♀; San Cristóbal de La Laguna; 28.4953, -16.3227; Aug. 2023; sus_scrofa, pers. obs. (<https://www.inaturalist.org/observations/187140681>) • 1 ♀; San Cristóbal de La Laguna; 28.5111, -16.3247; 21 Sep. 2023; Domingo Sosa, pers. obs. (<https://www.inaturalist.org/observations/184420785>) • 1 ♀; Santa Cruz de Tenerife; 28.4715, -16.2543; 3 Oct. 2023; D. Patiño, pers. obs. (<https://www.inaturalist.org/observations/186158783>) • 1 ♂; San Cristóbal de La Laguna; 28.4803, -16.3199; 12 Jun. 2024; nestork66,

pers. obs. (<https://www.inaturalist.org/observations/222221473>) • 1 ♀; Arafo, 28.3428, -16.4317; 13 Jul. 2024; wormsy, pers. obs. (<https://www.inaturalist.org/observations/228988344>) • 1 ♀; Santa Cruz de Tenerife; 28.4721, -16.2551; 11 Jul. 2024; skipshoemaker, pers. obs. (<https://www.inaturalist.org/observations/231008324>) • 1 ♀; Santa Cruz de Tenerife; 28.4639, -16.2730; 8 Sep. 2024; victor_febles, pers. obs. (<https://www.inaturalist.org/observations/240903947>). Spain. Canary Islands, **Gran Canaria**. 1 ♀; Maspalomas; 27.7420, -15.6000; 14 May 2023; alexsim123, pers. obs. (<https://www.inaturalist.org/observations/161581870>).

Discussion

Different observations obtained through citizen science in Gran Canaria and Tenerife, along with collected samples, confirm the presence of *A. florentinum* in the Canary Islands (Fig. 2).

Regarding the reproductive strategy of *A. florentinum*, resource defense polygyny is also displayed by males of sister species *A. manicatum* (Seidelmann 2021), arguably the most widespread unmanaged bee species globally (Strange et al. 2011). Different interactions between *A. manicatum* and native species of bees have been studied in regions beyond its native range, and some cases of resource competition (Soper and Beggs 2013) and exclusion (Graham et al. 2019) have been documented. Nonetheless, whether the arrival of *A. florentinum* to new territories poses the same threat for indigenous species is still uncertain, and further studies are needed to evaluate the impact of its introduction on native bee populations in the Canary Islands.

A. florentinum's generalist diet and its tendency to nest in pre-existing cavities enable this species to be easily transported for long distances, and eases its establishment in urban areas. Cities can host diverse communities of wild bee species, but they may also act as hotspots for biological invasions by allowing the accidental introduction of alien species, which in some cases leads to the biotic homogenization of urban communities of native bees, biodiversity loss, and the spread of exotic species into surrounding natural landscapes (Fitch et al. 2019). In this matter, the role of citizen science is crucial for the early detection and surveillance of invasive alien species, enabling public awareness on biological invasions and their negative impact on native biota (Roy et al. 2023).

Ever since the development of biodiversity monitoring platforms like iNaturalist and Observation.org, among others, volunteers can easily share georeferenced data of different taxa, based on observations, with the scientific community (Howard et al. 2022). Moreover, people with relevant expertise like taxonomists can also contribute by giving a more accurate identification to such records, therefore enhancing their taxonomic level and research quality (Callaghan et al. 2022). This is important because users that are less experienced at recognizing certain species or groups can sometimes fail in the identification process, as was the case with *A. florentinum* being initially mistaken for *A. manicatum*.




Figure 2. Localities where *A. florentinum* has been observed (red dots) in the islands of Tenerife and Gran Canaria.

This recent growing synergy between citizen scientists and experts has so far led to an increase of the knowledge about species' geographical distribution, and even to the rediscovery of species thought to be extinct or locally extirpated, new descriptions, and the early detection of alien species (Callaghan et al. 2022), that allows effective and cost-affordable fast management responses, as established populations are much more difficult and expensive to control (Roy et al. 2023). In the Canary Islands, the use of iNaturalist has proven to be useful for the discovery of new bee species such as the recently described *Pseudoanthidium jacobii* Vereecken & Litman, 2023, endemic to Fuerteventura and Lanzarote (Vereecken et al. 2023), and several exotic species besides *A. florentinum*, such as *Lasioglossum malachurum* (Kirby, 1802) in Tenerife (Domingo Sosa Obs. <https://www.inaturalist.org/observations/124150514>) and *Seladonia gemmea* (Dours, 1872) in Gran Canaria (biofreak Obs. <https://www.inaturalist.org/observations/137972057>).

Acknowledgements

We sincerely thank Gustavo Peña and David Lugo for their efforts in the collection and identification of specimens. We also thank Elizabeth Sauma and Antonio Mesa for their support in the sampling process. Special thanks to the reviewers and editor for the valuable feedback which contributed to improving this paper. We extend our sincere gratitude to Francisco La Roche for giving us access to his entomological

collection. Additionally, we would like to express our gratitude to the following iNaturalist users for generously sharing their observations: carmendgq, fer_guerra, Gustavo Peña, sus_scrofa, Domingo Sosa, nestork66, wormsy, skipshoemaker, victor_febles, and alexsim123.

References

- Aguib S, Louadi K, Schwarz M (2010) Les Anthidiini (Megachilidae, Megachilinae) d'Algérie avec trois espèces nouvelles pour ce pays: *Anthidium (Anthidium) florentinum* (Fabricius, 1775), *Anthidium (Proanthidium) amabile* Alfken, 1932 et *Pseudoanthidium (Exanthidium) enslini* (Alfken, 1928). Entomofauna 31: 121–152.
- Antropov AV, Astafurova YuV, Belokobylskij SA, Byvaltsev AM, Danilov YuN, Dubovikoff DA, et al. (2017) Annotated catalogue of the Hymenoptera of Russia. Volume I. Symphyta and Apocrita: Aculeata. Proceedings of the Zoological Institute. Russian Academy of Sciences. Supplement 6: 1–475. <https://doi.org/10.31610/trudyzin/2017.supl.6.5>
- Amiet F, Herrmann M, Müller A, Neumeyer R (2004) Apidae 4: *Anthidium*, *Chelostoma*, *Coelioxys*, *Dioxys*, *Heriades*, *Lithurgus*, *Megachile*, *Osmia*, *Stelis*. Fauna Helvetica 9: 1–273.
- Baldock DW, Livory A, Owens NW (2020) The bees and wasps of the Balearic Islands (Hymenoptera: Chrysidoidea, Vespoidea, Apoidea) with a discussion of aculeate diversity and endemism in Mediterranean and Atlantic archipelagos. Entomofauna, Supplement 25: 1–202.
- Balzan MV, Genoud D, Rasmont P, Schwarz M, Michez D (2017) New records of bees (Hymenoptera: Apoidea) from the Maltese Islands. Journal of Melittology 72: 1–9. <https://doi.org/10.17161/jom.v0i72.6626>
- Callaghan CT, Mesaglio T, Ascher JS, Brooks TM, Cabras AA, Chandler M, Cornwell WK, Cristóbal Ríos-Málaver I, Dankowicz E, Urfi Dhiya'ulhaq N, Fuller RA, Galindo-Leal C, Grattarola F, Hewitt S, Higgins L, Hitchcock C, James Hung KL, Iwane T, Kahumbu P, Kendrick R, Kieschnick SR, Kunz G, Lee CC, Lin CT, Loarie S, Norman Medina M, McGrouther MA, Miles L, Modi S, Nowak K, Oktaviani R, Waswala Olewe BM, Pagé J, Petrovan S, Saari C, Seltzer CE, Seregin AP, Sullivan JJ, Sumanapala AP, Takoukam A, Widness J, Willmott K, Wüster W, Young AN (2022) The benefits of contributing to the citizen science platform iNaturalist as an identifier. PLoS Biology 20(11): e3001843. <https://doi.org/10.1371/journal.pbio.3001843>
- Fitch G, Wilson CJ, Glaum P, Vaidya C, Simao MC, Jamieson MA (2019) Does urbanization favour exotic bee species? Implications for the conservation of native bees in cities. Biology Letters 15(12): 20190574. <https://doi.org/10.1098/rsbl.2019.0574>
- Fortunato L, Buian FM, Chiesa F, Zandigiacomo P (2013) Note biologiche su *Anthidium florentinum* nell'Italia nord-orientale (Hymenoptera, Megachilidae). Bollettino Società Naturalisti "Silvia Zenari", Pordenone 37: 137–145.
- GBIF (2024) GBIF. <https://www.gbif.org/> [Accessed 15 September 2024]
- Gogala A (2012) New records of bees (Hymenoptera: Apoidea: Anthophila) in Slovenia. Acta Entomologica Slovenica 20(1): 59–64.

- Graham KK, Eaton K, Obrien I, Starks PT (2019) *Anthidium manicatum*, an invasive bee, excludes a native bumble bee, *Bombus impatiens*, from floral resources. *Biological Invasions* 21: 1089–1099. <https://doi.org/10.1007/s10530-018-1889-7>
- Howard L, van Rees CB, Dahlquist Z, Luikart G, Hand BK (2022) A review of invasive species reporting apps for citizen science and opportunities for innovation. *NeoBiota* 71: 165–188. <https://doi.org/10.3897/neobiota.71.79597>
- iNaturalist (2024) iNaturalist. <https://www.inaturalist.org> [Accessed 15 September 2024]
- Kouakou D, Sattler T, Obrist MK, Duelli P, Moretti M (2008) Recent Swiss records of rare bee species (Hymenoptera, Apidae) with two species new to Switzerland. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 81: 191–197.
- Lieftinck MA (1958) A preliminary account of the bees of the Canary Islands (Hym., Apoidea). - *Commentationes biologicae* 18(5): 1–35.
- Lugo D, Peña G, De la Rúa P, Ruiz C (2022) Presence of exotic species of the wild bee genus *Hylaeus* (Hymenoptera: Colletidae) in the Canary Islands revealed by molecular and citizen science. *Journal of Apicultural Research* 61(4): 460–468. <https://doi.org/10.1080/00218839.2022.2053034>
- Niu ZQ, Yuan F, Ascher JS, Kasperek M, Orr MC, Griswold T, Zhu CD (2020) Bees of the genus *Anthidium* Fabricius, 1804 (Hymenoptera: Apoidea: Megachilidae: Anthidiini) from China. *Zootaxa* 4867(1): 1–67. <https://doi.org/10.11646/zootaxa.4867.1.1>
- Normandin É, Vereecken NJ, Buddle CM, Fournier V (2017) Taxonomic and functional trait diversity of wild bees in different urban settings. *PeerJ* 5: e3051. <https://doi.org/10.7717/peerj.3051>
- Ortiz-Sánchez FJ, La Roche F, Fuhrmann M (2016) Primera cita del género *Xylocopa* Latreille, 1802 en las Islas Canarias (Hymenoptera, Apidae). *Boletín de la Sociedad Entomológica Aragonesa* 58: 206.
- Romero D, Vargas P, Ornosca C (2021) On some features of the biology of the red snail-nesting bee, *Rhodanthidium sticticum* (Fabricius, 1787): phenology, flower preference, use of shells, flight ability and territorial behaviour. *Graellsia* 77(2): e146. <https://doi.org/10.3989/graellsia.2021.v77.307>
- Roy HE, Pauchard A, Stoett P, Renard Truong T, Bacher S, Galil B, Hulme PE, Ikeda T, Sankaran, KV, McGeoch, MA, Meyerson, LA, Nuñez M A, Ordonez A, Rahlao SJ, Schwindt E, Seebens H, Sheppard AW, Vandvik V (2023) Summary for Policymakers of the Thematic Assessment Report on Invasive Alien Species and their Control. IPBES Invasive Alien Species Assessment, 1–56. <https://doi.org/10.5281/zenodo.8314303>
- Ruiz C, Suárez D, Naranjo M, De la Rúa P (2020) First record of the carpenter bee *Xylocopa pubescens* (Hymenoptera, Apidae) in the Canary Islands confirmed by DNA barcoding. *Journal of Hymenoptera Research* 80: 169–175. <https://doi.org/10.3897/jhr.80.59649>
- Russo L (2016) Positive and negative impacts of non-native bee species around the world. *Insects* 7(4): 69. <https://doi.org/10.3390/insects7040069>
- Sardar S, Kasperek M, Rameshkumar A, Kazmi SI (2022) Four species of anthidiine bees (Apoidea: Megachilidae) new to India. *The Canadian Entomologist* 154(1): e7. <https://doi.org/10.4039/tce.2021.55>

- Schwenninger H (2008) Erster belegter Fund von *Anthidium florentinum* (Fabricius, 1775) in Deutschland (Hymenoptera: Apoidea: Megachilidae). *Mitteilungen des entomologischen Vereins Stuttgart* 43: 3–6.
- Seidelmann K (2021) Territoriality is just an option: allocation of a resource fundamental to the resource defense polygyny in the European wool carder bee, *Anthidium manicatum* (Hymenoptera: Megachilidae). *Behavioral Ecology and Sociobiology* 75(10): 138. <https://doi.org/10.1007/s00265-021-03073-6>
- Soper J, Beggs JR (2013) Assessing the impact of an introduced bee, *Anthidium manicatum*, on pollinator communities in New Zealand. *New Zealand Journal of Botany* 51(3): 213–228. <https://doi.org/10.1080/0028825X.2013.793202>
- Strange JP, Koch JB, Gonzalez VH, Nemelka L, Griswold T (2011) Global invasion by *Anthidium manicatum* (Linnaeus) (Hymenoptera: Megachilidae): assessing potential distribution in North America and beyond. *Biological Invasions* 13(9): 2115–2133. <https://doi.org/10.1007/s10530-011-0030-y>
- Strudwick T, Jacobi B (2018) The American Resin bee *Megachile (Chelostomoides) otomita* Cresson, 1878 established on Tenerife, Canary Islands (Spain) (Hymenoptera, Anthophila). *Ampulex* 10: 41–45.
- Sugiura S (2016) Impacts of introduced species on the biota of an oceanic archipelago: the relative importance of competitive and trophic interactions. *Ecological research* 31(2): 155–164. <https://doi.org/10.1007/s11284-016-1336-0>
- Varnava AI, Roberts SPM, Michez D, Ascher JS, Petanidou T, Dimitriou S, Devalez J, Pittara M, Stavrinides MC (2020) The wild bees (Hymenoptera, Apoidea) of the island of Cyprus. *ZooKeys* 924: 1–114. <https://doi.org/10.3897/zookeys.924.38328>
- Vereecken NJ, Ruiz C, Marshall L, Pérez-Gil M, Molenberg JM, Jacobi B, La Roche F, Litman JR (2023) A new small carder bee species from the eastern Canary Islands (Hymenoptera, Megachilidae, Anthidiini). *Journal of Hymenoptera Research* 96: 983–1015. <https://doi.org/10.3897/jhr.96.111550>
- Wirtz P, Kopka S, Schmoll G (1992) Phenology of two territorial solitary bees, *Anthidium manicatum* and *A. florentinum* (Hymenoptera: Megachilidae). *Journal of Zoology* 228(4): 641–651. <https://doi.org/10.1111/j.1469-7998.1992.tb04461.x>
- Zettel H, Zimmermann D, Wiesbauer H (2016) Ergänzungen zur Bienenfauna (Hymenoptera: Apidae) von Wien, Österreich. *Beiträge zur Entomofaunistik* 17: 85–107.